

- diagnosis and management of neuropathic pain. *Zhongguo Xian Dai Shen Jing Ji Bing Za Zhi*, 2010, 10:599-601. [李焰生.《神经病理性疼痛诊治专家共识》解读. 中国现代神经疾病杂志, 2010, 10:599-601.]
- [3] Attal N, Bouhassira D. Neuropathic pain: experimental advances and clinical applications. *Rev Neurol (Paris)*, 2004, 160:199-203.
- [4] Max MB. Clarifying the definition of neuropathic pain. *Pain*, 2002, 96:406-407.
- [5] Merskey H. Clarifying definition of neuropathic pain. *Pain*, 2002, 96:408-409.
- [6] Yu SY. Neuropathic pain. *Zhongguo Xian Dai Shen Jing Ji Bing Za Zhi*, 2010, 10:597-598. [于生元. 神经病理性疼痛. 中国现代神经疾病杂志, 2010, 10:597-598.]
- [7] Yu TM, Qi JJ. Clinical characteristics of neuropathic pain. *Zhongguo Xian Dai Shen Jing Ji Bing Za Zhi*, 2010, 10:608-610. [于挺敏, 齐晶晶. 神经病理性疼痛的临床特点. 中国现代神经疾病杂志, 2010, 10:608-610.]
- [8] Li F, Fang L, Huang S, Yang Z, Nandi J, Thomas S, Chen C, Camporesi E. Hyperbaric oxygenation therapy alleviates chronic constrictive injury-induced neuropathic pain and reduces tumor necrosis factor-alpha production. *Anesth Analg*, 2011, 113:626-633.
- [9] Zhai BZ, Ma Y, Tong XG. Efficacy observation of hyperbaric oxygen in the treatment of traumatic brain injury. *Zhongguo Xian Dai Shen Jing Ji Bing Za Zhi*, 2008, 8:161-163. [翟博智, 马越, 佟小光. 高压氧治疗颅脑创伤的疗效观察. 中国现代神经疾病杂志, 2008, 8:161-163.]
- [10] Peng Z, Wang S, Huang X, Xiao P. Effect of hyperbaric oxygen therapy on patients with herpes zoster. *Undersea Hyperb Med*, 2012, 39:1083-1087.
- [11] Kapoor S. Hyperbaric oxygen therapy and its emerging role in the management of non-neuropathic pain. *Eur J Pain*, 2012, 16:1204.
- [12] Hui J, Zhang ZJ, Zhang X, Shen Y, Gao YJ. Repetitive hyperbaric oxygen treatment attenuates complete freud's adjuvant - induced pain and reduces glia - mediated neuroinflammation in the spinal cord. *J Pain*, 2013, 14:747-758.
- [13] Thompson CD, Uhelski ML, Wilson JR, Fuchs PN. Hyperbaric oxygen treatment decreases pain in two nerve injury models. *Neurosci Res*, 2010, 66:279-283.
- [14] Bennett GJ, Xie YK. A peripheral mononeuropathy in rat that produces disorders of pain sensation like those seen in man. *Pain*, 1988, 33:87-107.
- [15] Chaplan SR, Bach FW, Pogrel JW, Chung JM, Yaksh TL. Quantitative assessment of tactile allodynia in the rat paw. *J Neurosci Methods*, 1994, 53:55-63.
- [16] Tsuda M, Shigemoto - Mogami Y, Koizumi S, Mizokoshi A, Kohsaka S, Salter MW, Inoue K. P2X4 receptors induced in spinal microglia gate tactile allodynia after nerve injury. *Nature*, 2003, 424:778-783.
- [17] Liu YQ. Progress of the treatment of neuropathic pain. *Zhongguo Xian Dai Shen Jing Ji Bing Za Zhi*, 2010, 10:611-614. [刘延青. 神经病理性疼痛治疗进展. 中国现代神经疾病杂志, 2010, 10:611-614.]
- [18] Xu H, Arita H, Hayashida M, Zhang L, Sekiyama H, Hanaoka K. Pain - relieving effects of processed Aconiti tuber in CCI - neuropathic rats. *J Ethnopharmacol*, 2006, 103:392-397.
- [19] Wang H, Li Y, Dun L, Xu Y, Jin S, Du J, Ma L, Li J, Zhou R, He X, Sun T, Yu J. Antinociceptive effects of oxymatrine from sophora flavescens, through regulation of NR2B - containing NMDA receptor - ERK/CREB signaling in a mice model of neuropathic pain. *Phytomedicine*, 2013, 20:1039-1045.
- [20] Gu N, Niu JY, Liu WT, Sun YY, Liu S, Lv Y, Dong HL, Song XJ, Xiong LZ. Hyperbaric oxygen therapy attenuates neuropathic hyperalgesia in rats and idiopathic trigeminal neuralgia in patients. *Eur J Pain*, 2012, 16:1094-1105.
- [21] Kroin JS, Takatori M, Li J, Chen EY, Buvanendran A, Tuman KJ. Upregulation of dorsal horn microglial cyclooxygenase - 1 and neuronal cyclooxygenase - 2 after thoracic deep muscle incisions in the rat. *Anesth Analg*, 2008, 106:1288-1295.
- [22] Patel RB, Pawar VD, Prajapati KD, Sonara BM, Deshpande SS, Shah GB, Jain MR. Anti-nociceptive and anti-allodynic activity of aliskiren in various pain models. *Eur J Pharmacol*, 2013, 708:80-87.
- [23] Shi XB, Yu SY. Pathogenesis of neuropathic pain. *Zhongguo Xian Dai Shen Jing Ji Bing Za Zhi*, 2010, 10:602-603. [时霄冰, 于生元. 神经病理性疼痛的发病机制. 中国现代神经疾病杂志, 2010, 10:602-603.]

(收稿日期:2013-08-09)

· 读者·作者·编者 ·

更正:脊髓小脑共济失调2型临床和神经影像学特征分析**Erratum to: Clinical and neuroimaging study of spinocerebellar ataxia type 2**

The article "Clinical and neuroimaging study of spinocerebellar ataxia type 2"^[1] published in Issue 6, Volume 13, 2013 has following corrections: 1) Page 525, Line 6 of Chinese Abstract; 2) Page 525, Line 10 of English Abstract; 3) Page 526, Line 19 on left column; 4) Page 528, Line 4 on right column.

"708例常染色体显性遗传性SCA2家系(708 autosomal dominant SCA2 families)" should be "708例常染色体显性遗传性SCA家系(708 autosomal dominant SCA families)".

Reference

- [1] Chen YY, Hao Y, Gu WH, Zhang J, Wang GX, Wang K, Jin M, Duan XH. Clinical and neuroimaging study of spinocerebellar ataxia type 2. *Zhongguo Xian Dai Shen Jing Ji Bing Za Zhi*, 2013, 13:525-532. [陈园园, 郝莹, 顾卫红, 张瑾, 王国相, 王康, 金森, 段晓慧. 脊髓小脑共济失调2型临床和神经影像学特征分析. 中国现代神经疾病杂志, 2013, 13:525-532.]